

POSTER PRESENTATION

Open Access

# Effect of post-exercise ingestion of different molecular weight carbohydrate solutions. Part III: Power output during a subsequent resistance training bout

Leighsa E Van Eck<sup>1\*</sup>, Anthony L Almada<sup>2</sup>, Margaret T Jones<sup>3</sup>, Andrew Jagim<sup>4</sup>, Joel Mitchell<sup>1</sup>, Jonathan M Oliver<sup>1</sup>

From The Twelfth International Society of Sports Nutrition (ISSN) Conference and Expo Austin, TX, USA. 11-13 June 2015

## Background

To maximize power adaptations, resistance training (RT) should be performed at maximal power output. In sports where more than one training bout is necessary in a day, subsequent RT may be limited by muscle glycogen, resulting in lower power output. High molecular weight (HMW) carbohydrate (CHO) solutions have been shown to result in greater glycogen re-synthesis rates, and greater work output during a subsequent cycling time trial compared to a low molecular weight (LMW) CHO solution. However, the effect of a HMW CHO on RT power output following exhaustive exercise is unknown.

## Methods

Sixteen resistance trained men (mean  $\pm$  SD; 23  $\pm$  3 years; 176.7  $\pm$  9.8 cm; 88.2  $\pm$  8.6 kg; 12.1  $\pm$  5.6% fat) participated in this study. One-repetition maximum (1RM) back squat (153.3  $\pm$  53.6 kg; 1.7  $\pm$  0.2 1RM:body mass), and  $\text{VO}_2$  max (37.4  $\pm$  4.3 ml·kg·min<sup>-1</sup>) were initially assessed in order to prescribe exercise intensities during experimental trials. In a double-blind, placebo-controlled, randomized cross over design consisting of three testing sessions separated by one week, subjects completed a glycogen depleting exercise bout on a cycle ergometer. Immediately post-exercise, subjects ingested a placebo (PLA), or a LMW or HMW CHO solution (10%) providing 1.2 kg·bw<sup>-1</sup> CHO, assigned randomly. Two hours post-ingestion, subjects performed 5 sets of 10 repetitions back squat (75% 1RM) "as explosively as possible". If subjects paused for more than 2 seconds or were unable to

complete a rep, resistance was lowered by 13.6 kg. Kinematic and kinetic measurements were sampled at 1000 Hz via force plate and two linear position transducers.

## Results

Average power following ingestion did not differ between CHO solutions until Set 4 ( $p = 0.108$ ) and Set 5 ( $p = 0.083$ ). Average power collapsed across the latter Sets was greater following ingestion of the HMW solution (Set 4, 1216  $\pm$  97 W; Set 5, 1143  $\pm$  102 W) compared to PLA (Set 4, 1066  $\pm$  80 W:  $p = 0.037$ ; Set 5, 1019  $\pm$  89 W:  $p = 0.048$ ), but not compared to ingestion of LMW (Set 4, 1160  $\pm$  79 W:  $p = 0.355$ ; Set 5, 1131  $\pm$  92 W:  $p = 0.852$ ). No difference was observed between LMW and PLA (Set 4,  $p = 0.275$ ; Set 5,  $p = 0.077$ ). The difference in average power was driven by velocity, as similar trends were observed in Set 4 and 5 ( $p = 0.100$  and  $p = 0.066$ , respectively). Average velocity was higher following ingestion of HMW (Set 4, 0.63  $\pm$  0.03 m·s<sup>-1</sup>; Set 5, 0.62  $\pm$  0.03 m·s<sup>-1</sup>) compared to PLA (Set 4, 0.56  $\pm$  0.04 m·s<sup>-1</sup>:  $p = 0.050$ ; Set 5, 0.56  $\pm$  0.04 m·s<sup>-1</sup>:  $p = 0.032$ ), but not LMW (Set 4, 0.61  $\pm$  0.03 m·s<sup>-1</sup>:  $p = 0.422$ ; Set 5, 0.61  $\pm$  0.03 m·s<sup>-1</sup>:  $p = 0.074$ ), with no difference between LMW and PLA (Set 4,  $p = 0.220$ ; Set 5,  $p = 0.769$ ). HMW conferred a likely beneficial effect in Sets 4 and 5 (92.5% and 88.7% likelihood, respectively), compared to PLA; while ingestion of LMW conferred only a possibly beneficial effect (68.7%) and likely beneficial effect (83.9%) in Sets 4 and 5, respectively.

## Conclusions

These data suggest post-exercise ingestion of a HMW CHO solution providing 1.2 kg·bw<sup>-1</sup> CHO may allow

<sup>1</sup>Department of Kinesiology, Texas Christian University, Fort Worth, TX 76129, USA

Full list of author information is available at the end of the article

athletes to sustain power output in a subsequent resistance training session when time between training sessions is limited.

#### Acknowledgements

This work was supported in part by Vitargo Global Sciences, LLC.

#### Authors' details

<sup>1</sup>Department of Kinesiology, Texas Christian University, Fort Worth, TX 76129, USA. <sup>2</sup>Vitargo Global Sciences, LLC, Dana Point, CA 92629, USA. <sup>3</sup>Health and Human Performance Division, George Mason University, Fairfax, VA 22030, USA. <sup>4</sup>Exercise & Sport Science Department, University of Wisconsin - La Crosse, La Crosse, WI 54601, USA.

Published: 21 September 2015

doi:10.1186/1550-2783-12-S1-P32

**Cite this article as:** Van Eck *et al.*: Effect of post-exercise ingestion of different molecular weight carbohydrate solutions. Part III: Power output during a subsequent resistance training bout. *Journal of the International Society of Sports Nutrition* 2015 **12**(Suppl 1):P32.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

